

COMPARISON OF SPHERICAL HARMONIC AND B-SPLINE MODELS FOR VTEC

MICHAEL SCHMIDT* AND DENISE DETTMERING*

*S German Geodetic Research Institute (DGFI)
Alfons-Goppel-Strasse 11 80539 Munich, Germany

Key words:

Abstract.

On the one-hand side modern satellite navigation and positioning systems require the use of precise and high resolution ionosphere correction models. On the other side the precise measurements of modern space-geodetic techniques, such as GNSS (terrestrial and space- borne) allow the study of ionosphere variations with an unprecedented accuracy. The characteristics of the ionosphere could be described by the electron density distribution and its functionals such as the vertical total electron content (VTEC).

Usually VTEC is modeled globally by a spherical harmonic expansion up to specific degree n (e.g. $n = 15$) for fixed time intervals (e.g. 2 hours). However, it is well-known that spherical harmonic models are not suited for representing data of heterogeneous density and quality. As a consequence, data gaps cannot be handled appropriately.

In this paper we present two different approaches for modeling VTEC generally depending on space and time:

1. The first and more traditional one is based on a system of 3-D base functions defined as the tensor product of spherical harmonics for longitude and latitude as well as endpoint-interpolating B-spline functions for the time.
2. In the second, alternative approach we define the 3-D system of base functions as tensor products of trigonometric B-spline functions for the longitude and two sets of endpoint-interpolating B-spline functions for latitude and time, respectively.

Whereas spherical harmonics are global base functions, the localizing feature of B-splines functions allows the use for global and regional applications. Thus, data gaps can be handled

appropriately. Furthermore, B-spline functions can be used to create a multi-scale analysis of VTEC, which can be used for

1. data compression and

2. near real-time processing using appropriate Kalman filtering.

We introduce both approaches mathematically and discuss advantages and disadvantages. In addition we apply the two methods and compare the results related to the evaluation of GNSS data.